


Sensory Systems Overview

From Sensation
to Perception...

Topics I	Topics II
Introduction & Electrochemical Gradients	Synaptic Transmission
Passive Membrane Properties	Electrophysiology Techniques
Action Potentials	Basic Circuits (Spinal Cord)
Voltage-Gated Ion Channels	Sensory Systems Overview
Ligand-Gated Ion Channels	Synaptic Plasticity

Study Material

- NEUROSCIENCE Third Edition
 - Dale Purves
- Unit II
 - Pages 189-337



THE COVER
Dorsal view of the human brain.
(Courtesy of S. Mark Williams.)

NEUROSCIENCE: Third Edition
Copyright © 2004 by Sinauer Associates, Inc. All rights reserved.
This book may not be reproduced in whole or in part without permission.

Address inquiries and orders to
Sinauer Associates, Inc.
23 Plumtree Road
Sunderland, MA 01375 U.S.A.

www.sinauer.com
FAX: 413-549-1118
orders@sinauer.com
publish@sinauer.com

Aims for this Lecture

- Know the main sensory systems and their transducers.
- Understand sensation as an active process.
We are not video cameras with smell detection tracks.....
- Learn about salient feature extraction along the processing pathway in different systems.


Recapitulation L8

- The stretch reflex response relies on one of the simplest complete functional circuits.
- We have put together the necessary elements and looked at their function.
- Muscle spindle, stretch receptor, afferent axon, synapse, motoneuron, efferent axon, neuromuscular endplate, muscle...

Mechanical stimuli and temperature


THE SOMATOSENSORY SYSTEM

The Sensors




Transducer Responses


Stimulus



Slowly adapting



Rapidly adapting



0

1


2

3

4

Time (s)

This an example of feature extraction at a very early stage. A change in the stimulus is more important than the constant presence of a stimulus.



Somatosensory Pathways

Information from the periphery (sensor) reaches the cortex in three synaptic steps.

This is the case for all senses except smell.

The information from all senses (except smell) passes through thalamic nuclei and is relayed there.

The thalamus is considered to be the gating structure for the flow of information to the cortex.


It receives many more top-down connections from the cortex than bottom-up connections from the periphery.

Light and dark

THE VISUAL SYSTEM

The Sensors

(A) Section of retina



(B)





Figure 10.4 Structure of the retina. (A) Section of the retina showing overall arrangement of retinal layers. (B) Diagram of the basic circuitry of the retina. A three-neuron chain—photoreceptor, bipolar cell, and ganglion cell—provides the most direct route for transmitting visual information to the brain. Horizontal cells and amacrine cells mediate lateral interactions in the outer and inner plexiform layers, respectively. The terms *inner* and *outer* designate relative distances from the center of the eye (inner, near the center of the eye; outer, away from the center, or toward the pigment epithelium).




Light travels through the inner layers of the retina to reach the photoreceptors.


The retina is the only part of the central nervous system that is directly visible from the outside

Visual System Transducer

► Response of Rods to Light



Copyright © 2001 by Allyn & Bacon



Note the enormous dynamic range





Figure 10.5 An intracellular recording from a single cone stimulated with different amounts of light (the cone has been taken from the turtle retina, which accounts for the relatively long time course of the response). Each trace represents the response to a brief flash that was varied in intensity. At the highest light levels, the response amplitude saturates (at about -65 mV). The hyperpolarizing response is characteristic of vertebrate photoreceptors; interestingly, some invertebrate photoreceptors depolarize in response to light. (After Schnapf and Baylor, 1987.)

Feature Extraction



Visual processing already starts in the retina.

Ganglion cells can have on-center or off-center receptive fields.

Central Vision Pathways




Figure 11.2 Central projections of retinal ganglion cells. Ganglion cell axons terminate in the lateral geniculate nucleus of the thalamus, the superior colliculus, the pretectum, and the hypothalamus. For clarity, only the crossing axons of the right eye are shown (view is looking up at the inferior surface of the brain).



Central Responses

(A) Experimental setup







Figure 11.9 Neurons in the primary visual cortex respond selectively to oriented edges. (A) An anesthetized animal is fitted with contact lenses to focus the eyes on a screen, where images can be projected; an extracellular electrode records the neuronal responses. (B) Neurons in the primary visual cortex typically respond vigorously to a bar of light oriented at a particular angle and weakly—or not at all—to other orientations.

The feature extraction continues....


Hermann Grid Illusion



Café Wall Illusion (M. Gregory)




Context is Everything..



Detecting volatile chemicals


SMELL

General Organization



The sensory neurons in the olfactory epithelium are constantly regenerated – thus maintaining function in a hostile environment.

Signal Transduction



A very large family of G-protein coupled receptors.... > 900 members in humans..

Odorant Coding

